



Gliricidia-based doubled up technology for improved crop production and agroecosystem resilience in Kongwa and Kiteto Districts

Kimaro, A.A. (ICRAF), Hafner, J. (ZALF/ICRAF), E. Swai (TARI-Hombolo) and Okori, P. (ICRISAT)

Challenges and Study objective

- ✓ Extensive grazing and the use of crop residues for cooking energy are major drivers of land degradation in semiarid sites.
- ✓ Poor soil health and unreliable precipitations limit crop production and increase susceptibility to climate change
- ✓ Low use of inputs, especially mineral fertilizer, by farmers to sustain crop production

Main study objective

- To evaluate productivity and resilience effects of Gliricidia-based intercropping systems in semiarid Tanzania
- To test cropping arrangements/sequences that respond to farm sizes and agroecologies and as well as integrating farmer production objectives

Introduced technologies

- (i) Gliricidia-maize intercropping
- (ii) Gliricidia-maize-pigeonepea intercropping (Doubled up legume)

Evidence

- Intercropping maize with *G. sepium* and pigeonpea improved crops (maize and pigeonpea) yield by up to 33%, besides wood and fodder supply for improved livestock nutrition (Fig. 1).
- Intercropping at the appropriate proportions based on local site conditions optimizes crops yield and builds agroecosystem resilience as higher legume proportions (1:2 ratio of maize to pigeonpea) was more beneficial in a less potential site (Mlali) than a higher potential site (Chitego)(Table 1).
- Increased economic benefits of the doubled up technology compared to sole maize as noted by 3- and 4- folds increase in return to labour per day worked and gross margin, respectively (Table 2)
- Environmental benefits include improved soil fertility and organic carbon (45%), rainwater use efficiency (37%) and offsetting CO₂ emissions through on-farm wood supply (Kimaro et al., 2007; Kimaro et al., 2016)

Approaches of scaling the technologies

- Since 2014 the project has trained 151 contact farmers (52.5% Female) on tree seedling production and agroforestry technologies for semiarid sites.
- Agricultural shows, field days and other dissemination approaches has been used to out scale the technology to over 6,000 farmers (Fig. 2).
- Partnered with district council, extension officers and contact farmers to out scale the technology to Ngumbi and Kilosa.

Partners:



We thank farmers and local partners in Africa RISING sites for their contributions to this work. We also acknowledge the support of all donors which globally support the work of the CGIAR centers and their partners through their contributions to the [CGIAR system](https://www.cgiar.org/)



This poster is licensed for use under the Creative Commons Attribution 4.0 International Licence. September 2018




Fig. 1: Maize (a) and pigeonpea (b) grain yields in the Doubled up legume systems at Manyusi, Kongwa. MMGlr=Maize+Gliricidia; MMMM=Maize monoculture, MMPP = Maize+Pigeonpea, MMPPGlr=Maize+Pigeonpea+Gliricidia, PPPP=Pure pigeonpea. Basal P was applied to all plots at planting at a rate of 15 kg P ha⁻¹ using Minjingu (NAFAKA) fertilizer.

Table 1: LER for Maize(MM)-pigeonpea (PP) intercropping at Mlali and Chitego Villages, Tanzania (Kimaro et al. 2018, In press)

Maize-PP ratio	2015		2016	
	Mlali	Chitego	Mlali	Chitego
2M:1PP	1.13	1.56	1.21	1.17
1M:1PP	1.12	1.47	1.46	1.53
1M:2PP	1.32	1.15	1.54	1.28



Fig. 2: Improved crops growth in a doubled up legume at Mlali Village (a). The Vice President of Zanzibar, Hon. Ambassador Sefu Ali Idi receiving information on the Gliricidia-maize intercropping technology at a demonstration plot in the agricultural exhibition ground in Morogoro in August 2018 (b).

Table 2: Gross Margin and Return to Labor for Maize (MM)-Gliricidia (GS)-pigeonpea (PP) technology at Manyusi Village

Maize-GS-PP Intercropping	2015		2016	
	GM (USD/ha)	RNLB	GM (USD/ha)	RNLB
MMPPGlr	487	11.4	1263	24.0
MMGlr	149	5.20	141	5.00
MMPP	1096	20.4	1715	30.4
MMMM	155	4.40	257	6.20
PPPP	1060	19.2	1669	29.0

Proposals for the future

- Evaluate drought resilience and trade-offs of the double up technology in semiarid.
- Assess soil health and mitigation co-benefits of the doubled up legume, which are just being realized.
- Capacity building and strengthening partnerships with national partners to supporting wider scaling of the technology

